



## Scientific note

# Tail bifurcation in *Tropidurus hispidus* (Squamata, Tropiduridae), in a semiarid area of Northeastern Brazil

Juliana Delfino de Sousa<sup>1</sup>, Maria Eduarda de Araújo Almeida<sup>1,2</sup>, José Henrique de Andrade Lima<sup>1,3,4</sup>, Alfredo dos Santos Almeida<sup>1</sup> & Marcelo Nogueira de Carvalho Kokubum<sup>1,2,5\*</sup>

*Tropidurus hispidus* is a member of the Tropiduridae family and presents a wide distribution in Northeastern Brazil (Abreu *et al.* 2002.). It is a diurnal, heliophilous, generalist, and abundant species (Vitt *et al.* 1996) and is mainly observed in rocky outcrops, trunks, and man-made structures (Rodrigues 1987, Carvalho 2013). Herein, we report two cases of tail bifurcation in *Tropidurus hispidus*.

On the 3rd September 2017 at 0839 h in an area of Caatinga, in a rural property named Sítio Fechado, (7° 20' 45" S/37° 17' 20" W; 737 m.a.s.l) in the municipality of Brejinho, Pernambuco State, Northeastern Brazil, we sighted an adult *T. hispidus* female presenting a bifurcated tail, on the wall of a residence (Figure 1). The second sighting was recorded on the 24th September 2019 at 1150h, during a behavioral study of this species (primary author), we observed one second female presenting a small tail bifurcation, during a mating encounter (Figure 2). Lizards have a remarkable ability to repair various tissues, such as muscle, osteological and

neuronal, in addition to their notable ability to regenerate their damaged tails (Alibardi 2010). Tail malformations can result from failures in regeneration after autotomy or may be congenital (Conzendey *et al.* 2013). It is considered an anti-predatory strategy, whose main objective is to increase the survival rate when the escape behavior is inefficient and may also be related to intraspecific aggression during mating and territoriality (Arnold 1994, Koleska *et al.* 2017, Bateman & Fleming 2019). In addition, it may be related to adverse environmental conditions caused by pollutants or pathogens that cause these malformations (Martinelli & Bogan 2013).

Some records of bifurcated tails have been highlighted in the available literature on the lizard species *Tropidurus torquatus* (Martins *et al.* 2013, Martinelli & Bogan 2013) *Tropidurus semitaeniatus* (Passos *et al.* 2014), *Vanzosaura rubricauda* (Pheasey *et al.* 2014), *Aspidoscelis tigris septentrionalis* (Heyborne *et al.* 2019), *Hemidactylus agrisus*

<sup>1</sup> Laboratório de Herpetologia, Unidade Acadêmica de Ciências Biológicas, Centro de Saúde e Tecnologia Rural, Universidade Federal de Campina Grande, Av. dos Universitários, s/n, Santa Cecília, 58700-970, Patos-PB, Brasil.

<sup>2</sup> Programa de Pós-graduação em Ciências Florestais, Centro de Saúde e Tecnologia Rural, Universidade Federal de Campina Grande, Av. dos Universitários, s/n, Santa Cecília, 58700-970, Patos-PB, Brasil.

<sup>3</sup> Programa de Pós-Graduação em Biologia Animal, Departamento de Zoologia, Centro de Biociências, Universidade Federal de Pernambuco, Av. Professor Moraes Rego, 1235, Cidade Universitária, 50670-901, Recife, Pernambuco, Brazil.

<sup>4</sup> Laboratório de Herpetologia, Departamento de Zoologia, Centro de Biociências, Universidade Federal de Pernambuco, Av. Professor Moraes Rego, 1235, Cidade Universitária, 50670-901, Recife, Pernambuco, Brazil.

<sup>5</sup> Programa de Pós-graduação em Ecologia e Conservação, Universidade Estadual da Paraíba, Bodocongó, 58429-500, Campina Grande, Paraíba, Brazil.

\* Corresponding author e-mail: [mnckkokubum@gmail.com](mailto:mnckkokubum@gmail.com)

(Andrade *et al.* 2015) *Ameiva ameiva* (Gogliath *et al.* 2012), *Salvator merianae* (Pelegrin & Leão 2016), *Ameivula ocellifera* (Sales & Freire 2019), and *Iguana iguana* (Lozano & Siro 2020), lepidosaurs (Barr *et al.* 2020). Here we observed another case for a *Tropidurus* species, likely due to the widespread distribution and high abundance of this species in Northeastern Brazil, as well as its frequent occurrence in anthropized environments. Although this phenomenon is common in nature, it likely does not affect the survival and reproductive success of individuals, as has been reported in the literature (Martins *et al.* 2013, Passos *et al.* 2014, Barr *et al.* 2020) and our observation, as in one of our reports, the individual was in the process of mating.



**Figure 1.** Tail bifurcation in *Tropidurus hispidus* (female individual) Sítio Fechado, in the municipality of Brejinho, Pernambuco. Photo: Sousa, J.D (2017).



**Figure 2.** Tail bifurcation in *Tropidurus hispidus* (female individual), during copulation behavior, Sítio Fechado, in the municipality of Brejinho, Pernambuco. Photo: Sousa, J.D (2019).

## REFERENCES

- Abreu, M.L., Frota, J.G. & Yuki, R.N. (2002). Geographic distribution, *Tropidurus hispidus*. *Herpetological Review* 33: 66.
- Alibardi, L. (2010). *Morphological and Cellular Aspects of Tail and Limb Regeneration in Lizards: A Model System with Implications for Tissue Regeneration in Mammals*. Advances in Anatomy, Embryology, and Cell Biology. Berlin, Germany, Springer-Verlag. 128pp.
- Andrade, M.J.M., Lopes, J.R.I., Sales, R.F.D. & Freire, F.M.X. (2015). *Hemidactylus agrius* (Country leaf-toed gecko): Polydactyly and tail bifurcation. *The Herpetological Bulletin* 13: 28-29.
- Arnold, E.N. (1994). *Caudal autotomy as a defense*. In: Gans C. and R. B. Huey (Eds), *Biology of the Reptilia*. Branta Book, Washington 235-274.
- Bateman, P.W. & Fleming, P.A. (2009). To cut a long tail short: a review of lizard caudal autotomy studies carried out over the last 20 years. *Journal of Zoology* 277: 1-14.
- Barr J.I., Somaweera R., Godfrey S.S., Gardner M.G. & Bateman P.W. (2020). When one tail isn't enough: abnormal caudal regeneration in lepidosaurs and its potential ecological impacts. *Biological Reviews*. 95:1479-1496.
- Carvalho, A.L.G. (2013). On the distribution and conservation of the South American lizard genus *Tropidurus* Wied-Neuwied, 1825 (Squamata: Tropiduridae). *Zootaxa* 3640: 42-56.
- Conzendey, P., Campos, A.P.S., Mollanna, F., Amorim, J.D.C.G. & Sousa, B.M. (2013). *Ophiodes striatus* (Striped Worm Lizard). Tail bifurcation. *Herpetological Review* 44: 45-146.
- Gogliath, M., Pereira, L.C.M., Nicola, P.A. & Ribeiro, L.B. (2012). *Ameiva ameiva* (Giant Ameiva). Bifurcation. *Herpetological Review* 43: 129.
- Heyborne, W.H., Gardner, C.E., Shipley, I.W & Andre, N.S. (2019). *Aspidoscelis tigris septentrionalis* (Plateau Tiger Whiptail). Supernumerary caudal anomaly. *Herpetological Review* 50: 569-570.
- Koleska, D., Svobodova, V., Husák, T., Kulma, M. & Jablonski, D. (2017). Tail bifurcation recorded in *Sauromalus ater*. *Herpetology Notes* 10: 363-364.

- Lozano, J.A. & Siro, D.P. (2020). Regenerate tail bifurcation in the Green Iguana (*Iguana iguana* Linnaeus, 1758). *Herpetology Notes* 13: 483-484.
- Martinelli, A.G. & Bogan, Y.S. (2013). Caudal pathology in *Tropidurus torquatus* (Iguania, Tropiduridae) from northeast Argentina. *Natural History* 3: 93-97.
- Martins, R.L., Peixoto, P.G., Fonseca, P.H.M., Martinelli, A.G., Silva, W.R. & Pelli, A. (2013). Abnormality in the tail of the collared lizard *Tropidurus* gr. *torquatus* (Iguania, Tropiduridae) from Uberaba city, Minas Gerais State, Brazil. *Herpetology Notes* 6: 369-371.
- Passos, D.C., Pinheiro, L.T., Galdino, C.A.B. & Rocha, C.F.D. (2014). *Tropidurus semitaeniatus* (Calango de lagedo). Tail bifurcation. *Herpetological Review* 45: 138.
- Pelegrin, N. & Leão, S.M. (2016). Injured *Salvator merianae* (Teiidae) regenerates six tails in central Argentina. *Cuadernos de Herpetología* 30: 21-23.
- Pheasey, H., Smith, P., Brouard, J.P. & Atkinson, K. (2014). *Vanzosaura rubricauda* (Red-tailed Vanzosaura). Bifurcation and trifurcation. *Herpetological Review* 45: 138-139.
- Rodrigues, M.T. (1987). Systematics, ecology and zoogeography of the *Tropidurus* of the *torquatus* group occurring south of Rio Amazonas (Sauridae, Iguanidae). *Arquivos de Zoologia de São Paulo* 31: 105-230.
- Sales, R.F.D. & Freire, E.M.X. (2019). *Ameivula ocellifera* (Spix's Whiptail lizard?). Tail bifurcation. *Herpetological Review* 50: 780.
- Vitt, L.J., Zani, P.A. & Caldwell, J.P. (1996). Behavioural ecology of *Tropidurus hispidus* on isolated rock outcrops in Amazonia. *Journal of Tropical Ecology* 12: 81-101.