



Scientific Note

Biodiversity in an urban swamp forest fragment in Ubatuba, São Paulo, Brazil

João Albert Bachur¹ Lucas Botelho¹ Edelcio Muscat¹

¹Projeto Dacnis, Estrada do Rio Escuro 4754, Sertão das Cotias, 11680-000, Ubatuba, SP, Brasil

*Corresponding author: bachurjoao@gmail.com

Received 07 February 2025 | Accepted 31 December 2025 | Published 06 January 2026

How to cite: Bachur, J. A., Botelho, L. & Muscat, E. (2025) Biodiversity in an urban swamp forest fragment in Ubatuba, São Paulo, Brazil *Heringeriana*, 20(1), e918072. doi.org/10.70782/heringeriana.v20i1.918072

Abstract: Urbanization promotes changes in land use and the fragmentation of natural habitats giving rise to forest fragments within the urban matrix. We assessed the avifauna and herpetofauna of a 2.93 ha urban fragment of *Tabebuia cassinoides* (Lam.) DC. swamp forest in the center of Ubatuba, SP, located 2.2 km from the Serra do Mar State Park. Between August 2022 and August 2023, we conducted active searches totaling 192 man-hours. We recorded 72 bird species and 15 herpetofauna species. The fragment acts as a breeding site for amphibians and serves as a refuge and stepping stone for the species sampled, reinforcing its importance for the conservation of urban biodiversity.

Keywords: Urban matrix; stepping stone; herpetofauna; avifauna; *Tabebuia cassinoides*.

Resumo: (Biodiversidade em um fragmento urbano de floresta paludosa em Ubatuba, São Paulo, Brasil.) A urbanização promove mudanças no uso do solo e a fragmentação de habitats naturais, originando fragmentos florestais inseridos na matriz urbana. Avaliamos a avifauna e a herpetofauna de um fragmento urbano de 2,93 ha de floresta paludosa de *Tabebuia cassinoides* (Lam.) DC. no centro de Ubatuba, SP, localizado a 2,2 km do Parque Estadual da Serra do Mar. Entre agosto de 2022 e agosto de 2023, realizamos buscas ativas totalizando 192 horas-homem. Registramos 72 espécies de aves e 15 espécies de herpetofauna. O fragmento atua como sítio reprodutivo para anfíbios e funciona como refúgio e "stepping stone" das espécies amostradas, reforçando sua importância para a conservação da biodiversidade urbana.

Palavras-chave: Matriz urbana; "stepping stone"; herpetofauna; avifauna; *Tabebuia cassinoides*.

Urbanization comes with changes in the environmental use of soil, as well as natural habitat fragmentation (Haddad et al., 2015; Liu et al., 2016), giving rise to the concept of urban forests (Giacon et al., 2022). These forest fragments become green islands in a complex urban matrix. They are fundamental for biodiversity and provide critical habitat for a variety of species, as well as key ecosystem services for human populations (Canedoli et al., 2018; Long & Frank, 2020; Giaccon et al., 2022; Mitchell & Devisscher, 2022; Ribeiro et al., 2022).

Although they are isolated, urban forest fragments can be important connectors with protected areas nearby, making genetic flow possible between them as observed in different studies around the world (Parris, 2006; Van Rossum & Triest, 2012). The animal, seed, and pollen dispersion through fragments and protected areas minimize the adverse effects of habitat fragmentation on biodiversity, such as diversity loss (Antonelli et al., 2022; Ribeiro et al., 2022). Moreover, those fragments are also an ex-

tension of species' home ranges, amplifying their demography.

In the center of Ubatuba - São Paulo there is a 2.93 hectare fragment of *Tabebuia cassinoides* (Lam.) DC. (Bignoniaceae) swamp forest. It belongs to the town administration and is adjacent to the local military police battalion, the fire department, and the municipal convention center. It is 2.2 kilometers in a straight line from the closest part of the Serra do Mar State Park, coordinates -23.444098 S, -45.071359 O, 3 m above sea level (Figures 1A and 1B). The fragment is surrounded by urban landscape, and part of it floods seasonally. Its vegetation and terrestrial fauna are isolated, but it functions as a refuge. *Tabebuia cassinoides* (Lam.) DC. is an endangered species, VU (vulnerable), according to the Brazilian Flora Red List (Martinelli & Moraes, 2013), and few regions still have swamp forests. Studying this urban forest could yield important information about the biodiversity of a threatened ecosystem inserted in a city.




Figure 1: **1A.** Urban fragment studied (orange) and the Serra do Mar State Park boundaries (bright green), which are 2.2 km away from the fragment in a straight line; **1B.** Close-up of the study area in the Ubatuba city center.

We monitored the swamp forest fragment from August 2022 to August 2023, concentrating on avifauna and herpetofauna. Two researchers conducted active searches

twice a month: one nocturnal from 19:00 to 21:00, focusing more on amphibians and reptiles; and one diurnal between 06:00 and 08:00, looking for birds, adding up to 192

man-hours. The active search was conducted within the borders of the fragment, especially around three ponds in the area. There were no marked trails. All species of avifauna and herpetofauna recorded during the searches were entered into a database, including birds found at night and herpetofauna seen during the day.

We recorded 72 bird species from 15 different orders, 30 families and 68 genera (Table 1). The highlights

of the list were the Little Nightjar *Hydropsalis parvula* (Gould, 1837) (Figure 2F) and the Rusty-backed Spine-tail *Cranioleuca vulpina* (Pelzeln, 1856). Both species are rarely present in urban areas and were absent in previous large-scale bird surveys in Ubatuba (Dario & De-Vincenzo, 2011; Simpson et al., 2012). There are only a few records for the municipality in the digital citizen science platforms WikiAves and eBird (eBird, 2021; WikiAves, 2024b; WikiAves, 2024a).

Table 1. List of bird species recorded during the monitoring of the urban forest fragment in Ubatuba, SP, as well as the conservation status in the São Paulo state registries (1) and in the national (2) (São Paulo, 2018; ICMBio, 2025) (* = migratory species).

ORDER	Family	Scientific name	Popular name	1	2
ANSERIFORMES	Anatidae	<i>Amazonetta brasiliensis</i> (Gmelin, JF, 1789)	Brazilian Teal*	LC	LC
COLUMBIFORMES	Columbidae	<i>Patagioenas picazuro</i> (Temminck, 1813)	Picazuro, Pigeon	LC	LC
		<i>Columbina talpacoti</i> (Temminck, 1810)	Ruddy Ground, Dove	LC	LC
		<i>Columba livia</i> Gmelin, JF, 1789	Rock Dove	X*	NE
NYCTIBIIFORMES	Nyctibiidae	<i>Nyctibius griseus</i> (Gmelin, JF, 1789)	Common Potoo	LC	LC
CAPRIMULGIFORMES	Caprimulgidae	<i>Hydropsalis parvula</i> (Gould, 1837)	Little Nightjar	LC	LC
GRUIFORMES	Rallidae	<i>Aramides saracura</i> (Spix, 1825)	Slaty-breasted, Wood Rail	LC	LC
APODIFORMES	Apodidae	<i>Chaetura meridionalis</i> Hellmayr, 1907	Sick's Swift	LC	LC
		<i>Progne chalybea</i> (Gmelin, JF, 1789)	Grey-breasted Martin	LC	LC
		<i>Pygochelidon cyanoleuca</i> (Vieillot, 1817)	Blue-and-white Swallow	LC	LC
	Trochilidae	<i>Chionomesa fimbriata</i> (Gmelin, JF, 1788)	Glittering-throated Emerald*	LC	LC
		<i>Florisuga fusca</i> (Vieillot, 1817)	Black Jacobin*	LC	LC
		<i>Eupetomena macroura</i> (Gmelin, JF, 1788)	Swallow-tailed Hummingbird	LC	LC
		<i>Thalurania glaucopis</i> (Gmelin, JF, 1788)	Violet-capped Woodnymph	LC	LC
		<i>Vanellus chilensis</i> (Molina, 1782)	Southern Lapwing	LC	LC
		<i>Mesembrinibis cayennensis</i> (Gmelin, JF, 1789)	Green Ibis	LC	LC
		<i>Phimosus infuscatus</i> (Lichtenstein, MHC, 1823)	Bare-faced Ibis	LC	LC
CHARADRIIFORMES	Threskiornithidae	<i>Syrigma sibilatrix</i> (Temminck, 1824)	Whistling Heron	LC	LC
		<i>Butorides striata</i> (Linnaeus, 1758)	Striated Heron	LC	LC
		<i>Ardea alba</i> Linnaeus, 1758	Great Egret	LC	LC
	Phalacrocoracidae	<i>Nannopterum brasilianum</i> (Gmelin, JF, 1789)	Neotropic Cormorant	LC	LC
		<i>Coragyps atratus</i> (Bechstein, 1793)	Black Vulture	LC	LC
		<i>Rupornis magnirostris</i> (Gmelin, JF, 1788)	Roadside Hawk	LC	LC
		<i>Asio clamator</i> (Vieillot, 1808)	Striped Owl	LC	LC
		<i>Colaptes campestris</i> (Vieillot, 1818)	Campo Flicker	LC	LC
		<i>Picumnus cirratus</i> Temminck, 1825	White-barred Piculet	LC	LC
		<i>Milvago chimachima</i> (Vieillot, 1816)	Yellow-headed, Caracara	LC	LC
FALCONIFORMES	Falconidae	<i>Caracara plancus</i> (Miller, JF, 1777)	Crested Caracara	LC	LC
		<i>Falco femoralis</i> Temminck, 1822	Aplomado Falcon	LC	LC
		<i>Brotogeris tirica</i> (Gmelin, JF, 1788)	Plain Parakeet	LC	LC
	Psittacidae	<i>Pyrrhura frontalis</i> (Vieillot, 1818)	Maroon-bellied Parakeet	LC	LC
		<i>Forpus xanthopterygius</i> (Spix, 1824)	Cobalt-rumped Parrotlet	LC	LC
		<i>Herpsilochmus rufimarginatus</i> (Temminck, 1822)	Rufous-margined Antwren	LC	LC
		<i>Dysithamnus mentalis</i> (Temminck, 1823)	Plain Antwren	LC	LC
		<i>Lepidocolaptes angustirostris</i> (Vieillot, 1818)	Narrow-billed Woodcreeper	LC	LC
		<i>Furnarius figulus</i> (Lichtenstein, MHC, 1823)	Band-tailed Hornero	LC	LC
		<i>Synallaxis spixi</i> Sclater, PL, 1856	Spix's Spinetail	LC	LC
PASSERIFORMES	Thamnophilidae	<i>Cranioleuca vulpina</i> (Pelzeln, 1856)	Rusty-backed Spinetail*	LC	LC
		<i>Furnarius rufus</i> (Gmelin, JF, 1788)	Rufous Hornero	LC	LC
		<i>Todirostrum cinereum</i> (Linnaeus, 1766)	Common Tody-Flycatcher	LC	LC
	Dendrocolaptidae	<i>Tangus sulphuratus</i> (Linnaeus, 1766)	Great Kiskadee	LC	LC
		<i>Myiozetetes similis</i> (Spix, 1825)	Social Flycatcher	LC	LC
		<i>Elaenia flavogaster</i> (Thunberg, 1822)	Yellow-bellied Elaenia	LC	LC
	Furnariidae	<i>Cnemotriccus fuscatus</i> (Wied-Neuwied, M, 1831)	Fuscous Flycatcher	LC	LC
		<i>Pyrocephalus rubinus</i> (Boddaert, 1783)	Scarlet Flycatcher*	LC	LC
		<i>Campostoma obsoletum</i> (Temminck, 1824)	Southern Beardless Tyrannulet	LC	LC
		<i>Megarynchus pitangua</i> (Linnaeus, 1766)	Boat-billed Flycatcher	LC	LC
		<i>Tyrannus melancholicus</i> Vieillot, 1819	Tropical Kingbird*	LC	LC
		<i>Lathrotriccus eulerei</i> (Cabanis, 1868)	Euler's Flycatcher*	LC	LC
		<i>Fluvicola nengeta</i> (Linnaeus, 1766)	Masked Water Tyrant	LC	LC
		<i>Myiarchus ferox</i> (Gmelin, JF, 1789)	Short-crested Flycatcher	LC	LC
		<i>Tyrannus savana</i> Daudin, 1802	Fork-tailed Flycatcher*	LC	LC

Cont.

ORDER	Family	Scientific name	Popular name	1	2
PASSERIFORMES	Troglodytidae	<i>Troglodytes musculus</i> Naumann, JF, 1823	Southern House Wren	LC	LC
		<i>Cantorchilus longirostris</i> (Vieillot, 1819)	Long-billed Wren	LC	LC
	Turdidae	<i>Turdus albicollis</i> Vieillot, 1818	White-necked Thrush	LC	LC
		<i>Turdus leucomelas</i> Vieillot, 1818	Pale-breasted Thrush	LC	LC
		<i>Turdus rufiventris</i> Vieillot, 1818	Rufous-bellied Thrush	LC	LC
		<i>Turdus amaurochalinus</i> Cabanis, 1851	Creamy-bellied Thrush	LC	LC
		<i>Passer domesticus</i> (Linnaeus, 1758)	House Sparrow	X*	NE
	Fringillidae	<i>Euphonia pectoralis</i> (Latham, 1801)	Chestnut-bellied Euphonia	LC	LC
	Passerellidae	<i>Zonotrichia capensis</i> (Müller, PLS, 1776)	Rufous-collared Sparrow	LC	LC
	Icteridae	<i>Gnorimopsar chopi</i> (Vieillot, 1819)	Chopi Blackbird	LC	LC
		<i>Molothrus bonariensis</i> (Gmelin, JF, 1789)	Shiny Cowbird	LC	LC
	Parulidae	<i>Geothlypis aequinoctialis</i> (Gmelin, JF, 1789)	Masked Yellowthroat	LC	LC
	Thraupidae	<i>Ramphocelus bresilia</i> (Linnaeus, 1766)	Brazilian Tanager	LC	LC
		<i>Tangara seledon</i> (Müller, PLS, 1776)	Green-headed Tanager	LC	LC
		<i>Tachyphonus coronatus</i> (Vieillot, 1822)	Ruby-crowned Tanager	LC	LC
		<i>Coereba flaveola</i> (Linnaeus, 1758)	Bananaquit	LC	LC
		<i>Thraupis palmarum</i> (Wied-Neuwied, M, 1821)	Palm Tanager	LC	LC
		<i>Sicalis flaveola</i> (Linnaeus, 1766)	Saffron Finch	LC	LC
		<i>Thraupis sayaca</i> (Linnaeus, 1766)	Sayaca Tanager	LC	LC
		<i>Volatinia jacarina</i> (Linnaeus, 1766)	Blue-black Grassquit	LC	LC




Figure 2: Pictures of biodiversity and environment in the monitored swamp forest. A. Flooded environment dominated by caixeta (*Tabebuia cassinoides*). B. Orchid *Cattleya intermedia*. C. Snake Brazilian Sipo (*Chironius laevicolis*). D. Amplexus of Rio Mutum Snouted Tree Frog (*Ololygon argyreonatus*). E. White-headed Whistling Duck (*Dendrocygna viduata*). F. Little Nightjar (*Hydrosalis parvula*), an uncommon bird in Ubatuba, SP, photographed in the study area. Author: Projeto Dacnis collection.

Regarding herpetofauna, we found 13 amphibians of three families and eight genera, and two snakes from different families (Table 2). We observed that the forest fragment functions as a reproductive site for the frog species encountered, as at times there was intense vocalization from most of the recorded species (Figure 2D; Figure 3).

It shows that the fragment is a habitat for the metapopulation that occurs in the area. Moreover, although fragmentation and isolation of populations may result in genetic erosion and low variability, studies have shown that some species maintained high genetic diversity despite habitat fragmentation (Ralimanana et al., 2022).

Knowing the biodiversity of this urban fragment is fundamental to reinforce the need for its conservation. Since it lies less than 3 km away from protected forest in the Serra do Mar State Park, it may work as a stepping stone for local fauna populations, especially birds,

becoming a part of their home range (Galina & Gimenes, 2006). Forest fragments can also hold a high genetic diversity as well as extend the range of animal species (Tee et al., 2018; Ribeiro et al., 2022; Ralimanana et al., 2022).

Table 2. List of amphibians and reptiles recorded during the monitoring of the urban forest fragment in Ubatuba, SP, as well as the conservation status in the São Paulo state registries (1) and in the national (2) (São Paulo, 2018; ICMBio, 2025 (* = migratory species)).

ORDER	Family	Scientific name	Popular name	1	2
ANURA	Hylidae	<i>Boana albomarginata</i> (Spix, 1824)	White-banded Tree Frog	LC	LC
		<i>Boana faber</i> (Wied-Neuwied, 1821)	Smith Frog	LC	LC
		<i>Dendropsophus elegans</i> (Wied-Neuwied, 1824)	Elegant Forest Treefrog	LC	LC
		<i>Dendropsophus minutus</i> (Peters, 1872)	Lesser Treefrog	LC	LC
		<i>Itapotihyla langsdorffii</i> (Duméril and Bibron, 1841)	Ocellated Treefrog	LC	LC
		<i>Ololygon argyreomata</i> (Miranda-Ribeiro, 1926)	Rio Mutum Snouted Treefrog	LC	LC
		<i>Ololygon littoralis</i> (Pombal and Gordo, 1991)	Rio Verde Snouted Treefrog	LC	LC
		<i>Ololygon faivovichii</i> (Brasileiro, Oyamauchi, and Haddad, 2007)	Bandeirantes Snouted Treefrog	LC	LC
		<i>Scinax hayii</i> (Barbour, 1909)	Hay's Snouted Treefrog	LC	LC
		<i>Scinax imbegue</i> Nunes, Kwet, and Pombal, 2012	Snouted Treefrog	LC	LC
		<i>Trachycephalus mesophaeus</i> (Hensel, 1867)	Porto Alegre Golden-eyed Treefrog	LC	LC
	Leptodactylidae	<i>Leptodactylus latrans</i> (Steffen, 1815)	Argus Frog	LC	LC
	Phyllomedusidae	<i>Pithecopus rohdei</i> (Mertens, 1926)	Rohde's Frog	LC	LC
	Colubridae	<i>Chironius laevicollis</i> Wied-Neuwied, 1824	Brazilian sipo	LC	LC
	Dipsadidae	<i>Erythrolamprus miliaris</i> Linnaeus, 1758	Military Ground Snake	LC	LC
SQUAMATA					



Figure 3: Rhode's Frog (*Pithecopus rohdei*) vocalizing upside down in the area. Author: Projeto Dacnis collection.

Impacts of the fragmentation of forests, such as patch isolation and increased edge effects, lead to loss of biodiversity, due to the decrease in landscape connectivity. We need to improve our knowledge about the role of fragments in maintaining biodiversity in the urban matrix and of fauna distribution between them (Tee et al., 2018).

On the conservation side, environmental education initiatives involving communities close to forest fragments should raise awareness about their importance and reinforce the need for the survival of native urban green areas. Besides providing shelter for wildlife in an otherwise hostile setting, these areas are crucial to the maintenance of local biodiversity, as has been proven by the wide variety of fauna found in this small forest fragment in the Ubatuba city center.

Acknowledgments

We thank Elsie L. Rotenberg for her careful revision of the text.

References

- Antonelli, A., Smith, R. J., Perrigo, A. L., Crottini, A., Hackel, J., Testo, W., Farooq, H., Torres Jiménez, M. F., Andela, N., & Andermann, T. (2022). Madagascar's extraordinary biodiversity: Evolution, distribution, and use. *Science*, 378(6623), eabf0869. <https://doi.org/10.1126/science.abf0869>
- Canedoli, C., Manenti, R., & Padoa-Schioppa, E. (2018). Birds biodiversity in urban and periurban forests: environmental determinants at local and landscape scales. *Urban Ecosystems*, 21(4), 779–793. <https://doi.org/10.1007/s11252-018-0757-7>
- Dario, F. R., & De-Vincenzo, M. C. V. (2011). Avian diversity and relative abundance in a restinga forest of São Paulo, Brazil. *Tropical Ecology*, 52(1), 25–33.
- eBird. (2021). eBird: An online database of bird distribution and abundance. <http://www.ebird.org/>
- Galina, A. B., & Gimenes, M. R. (2006). Riqueza, composição e distribuição espacial da comunidade de aves em um fragmento florestal urbano em Maringá, Norte do Estado do Paraná, Brasil. *Acta Scientiarum. Biological Sciences*, 28(4), 379–388.
- Giacon, V. P., Valente, R. A., & Cardoso-Leite, E. (2022). Association between urbanization and the biotic integrity of urban forest remnants. *Ambiente & Sociedade*, 25, e00534. <https://doi.org/10.1590/1809-4422asoc20200053r4OA>
- Haddad, N. M., Brudvig, L. A., Clobert, J., Davies, K. F., Gonzalez, A., Holt, R. D., Lovejoy, T. E., Sexton, J. O., Austin, M. P., & Collins, C. D. (2015). Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science Advances*, 1(2), e1500052. <https://doi.org/10.1126/sciadv.1500052>
- ICMBio. (2025). Sistema de Avaliação do Risco de Extinção da Biodiversidade – SALVE. <https://salve.icmbio.gov.br/>
- Liu, Z., He, C., & Wu, J. (2016). The relationship between habitat loss and fragmentation during urbanization: An empirical evaluation from 16 world cities. *PloS one*, 11(4), e0154613. <https://doi.org/10.1371/journal.pone.0154613>
- Long, L. C., & Frank, S. D. (2020). Risk of bird predation and defoliating insect abundance are greater in urban forest fragments than street trees. *Urban Ecosystems*, 23(3), 519–531. <https://doi.org/10.1007/s11252-020-00939-x>
- Martinelli, G., & Moraes, M. A. (2013). Livro Vermelho da Flora do Brasil, 1100.
- Mitchell, M. G., & Devisscher, T. (2022). Strong relationships between urbanization, landscape structure, and ecosystem service multifunctionality in urban forest fragments. *Landscape and Urban Planning*, 228, e104548. <https://doi.org/10.1016/j.landurbplan.2022.104548>
- Parris, K. M. (2006). Urban amphibian assemblages as metacommunities. *Journal of Animal Ecology*, 75(3), 757–764. <https://doi.org/10.1111/j.1365-2656.2006.01096.x>
- Ralimanana, H., Perrigo, A. L., Smith, R. J., Borrell, J. S., Faurby, S., Rajaonah, M. T., Randriamboavonjy, T., Vorontsova, M. S., Cooke, R. S., & Phelps, L. N. (2022). Madagascar's extraordinary biodiversity: Threats and Opportunities. *Science*, 378(6623), eadf1466. <https://doi.org/10.1126/science.adf1466>
- Ribeiro, M. P., Mello, K., & Valente, R. A. (2022). How can forest fragments support protected areas connectivity in an urban landscape in Brazil? *Urban Forestry & Urban Greening*, 74, 127683. <https://doi.org/10.1016/j.ufug.2022.127683>
- São Paulo. (2018). Decreto nº 63.853 de 27 de novembro de 2018. <https://www.al.sp.gov.br/repositorio/legislacao/decreto/2018/decreto-63853-27.11.2018.html>
- Simpson, R., Cavarzere, V., & Simpson, E. (2012). List of documented bird species from the municipality of Ubatuba, state of São Paulo, Brazil. *Papéis Avulsos de Zoologia*, 52, 233–254. <https://doi.org/10.1590/S0031-10492012002100001>
- Tee, S. L., Samantha, L. D., Kamarudin, N., Akbar, Z., Lechner, A. M., Ashton-Butt, A., & Azhar, B. (2018). Urban forest fragmentation impoverishes native mammalian biodiversity in the tropics. *Ecology and Evolution*, 8(24), 12506–12521. <https://doi.org/10.1002/ece3.4632>
- Van Rossum, F., & Triest, L. (2012). Stepping-stone populations in linear landscape elements increase pollen dispersal between urban forest fragments. *Plant Ecology and Evolution*, 145(3), 332–340. <https://doi.org/10.5091/plecevo.2012.737>

WikiAves. (2024a). *Mapa de registros da espécie arredio-do-rio (Cranioleuca vulpina)*. https://www.wikiaves.com.br/mapaRegistros_arredio-do-rio/

WikiAves. (2024b). *Mapa de registros da espécie bacurau-chitã (Hydropsalis parvula)*. https://www.wikiaves.com.br/mapaRegistros_bacurau-chinta/



This is an open-access article distributed under the terms of the Creative Commons Attribution License.